Table 1 Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2). School entry and adolescent vaccine age groups are shaded in gray.



Table 2 Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 month Behind, United States, 2021

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. **Always use this table in conjunction with Table 1 and the notes that follow.**

Children age 4 months through 6 years

Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	**4 weeks**	**8 weeks *and* at least 16 weeks after first dose.** Minimum age for the final dose is 24 weeks.		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days.	**4 weeks**	**4 weeks** Maximum age for final dose is 8 months, 0 days.		
Diphtheria, tetanus, and acellular pertussis	6 weeks	**4 weeks**	**4 weeks**	**6 months**	**6 months**
Haemophilus influenzae type b	6 weeks	**No further doses needed** if first dose was administered at age 15 months or older. **4 weeks** if first dose was administered before the 1st birthday. **8 weeks (as final dose)** if first dose was administered at age 12 through 14 months.	**No further doses needed** if previous dose was administered at age 15 months or older. **4 weeks** if current age is younger than 12 months ***and*** first dose was administered at younger than age 7 months ***and*** at least 1 previous dose was PRP-T (ActHib, Pentacel, Hiberix) or unknown. **8 weeks *and* age 12 through 59 months (as final dose)** if current age is younger than 12 months ***and*** first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months ***and*** first dose was administered before the 1st birthday ***and*** second dose was administered at younger than 15 months; OR if both doses were PRP-OMP (PedvaxHIB, Comvax) ***and*** were administered before the 1st birthday.	**8 weeks (as final dose)** This dose only necessary for children age 12 through 59 months who received 3 doses before the 1st birthday.	
Pneumococcal conjugate	6 weeks	**No further doses needed** for healthy children if first dose was administered at age 24 months or older. **4 weeks** if first dose was administered before the 1st birthday. **8 weeks (as final dose for healthy children)** if first dose was administered at the 1st birthday or after.	**No further doses needed** for healthy children if previous dose was administered at age 24 months or older. **4 weeks** if current age is younger than 12 months and previous dose was administered at <7 months old. **8 weeks (as final dose for healthy children)** if previous dose was administered between 7–11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was administered before age 12 months.	**8 weeks (as final dose)** This dose only necessary for children age 12 through 59 months who received 3 doses before age 12 months or for children at high risk who received 3 doses at any age.	
Inactivated poliovirus	6 weeks	**4 weeks**	**4 weeks** if current age is <4 years. **6 months (as final dose)** if current age is 4 years or older.	**6 months** (minimum age 4 years for final dose).	
Measles, mumps, rubella	12 months	**4 weeks**			
Varicella	12 months	**3 months**			
Hepatitis A	12 months	**6 months**			
Meningococcal ACWY	2 months MenACWY-CRM 9 months MenACWY-D 2 years MenACWY-TT	**8 weeks**	See Notes	See Notes	

Children and adolescents age 7 through 18 years

Meningococcal ACWY	Not applicable (N/A)	**8 weeks**			
Tetanus, diphtheria; tetanus, diphtheria, and acellular pertussis	7 years	**4 weeks**	**4 weeks** if first dose of DTaP/DT was administered before the 1st birthday. **6 months (as final dose)** if first dose of DTaP/DT or Tdap/Td was administered at or after the 1st birthday.	**6 months** if first dose of DTaP/DT was administered before the 1st birthday.	
Human papillomavirus	9 years	**Routine dosing intervals are recommended.**			
Hepatitis A	N/A	**6 months**			
Hepatitis B	N/A	**4 weeks**	**8 weeks *and* at least 16 weeks after first dose.**		
Inactivated poliovirus	N/A	**4 weeks**	**6 months** A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years or if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	**4 weeks**			
Varicella	N/A	**3 months** if younger than age 13 years. **4 weeks** if age 13 years or older.			